

EXPLICIT BOUNDS FOR LARGE GAPS BETWEEN SQUAREFREE INTEGERS

ANGEL KUMCHEV, WADE MCCORMICK, NATHAN MCNEU, ARIANA PARK,
RUSSELL SCHERR, AND WILLOW ZIEHR

ABSTRACT. We obtain explicit forms of the current best known asymptotic upper bounds for gaps between squarefree integers. In particular we show, for any $x \geq 2$, that every interval of the form $(x, x + 11x^{1/5} \log x]$ contains a squarefree integer. The constant 11 can be improved further, if x is assumed to be larger than a (very) large constant.

1. INTRODUCTION

An integer n is called *squarefree* if it is not divisible by the square of any prime p . More generally, if $k \geq 2$, n is called *k -free* if it is not divisible by p^k for any prime p ; 3-free integers, in particular, are also known as *cubefree*.

The asymptotic distribution of the k -free integers has been studied systematically, at least since the early 1900s, with a special focus on the squarefree case. Let $Q_k(x)$ denote the counting function of the k -free numbers up to x , and consider the error term $E_k(x)$ in the asymptotic formula

$$Q_k(x) = \frac{x}{\zeta(k)} + E_k(x),$$

where $\zeta(k)$ is the Riemann zeta-function. The bound $E_k(x) = O(x^{1/k})$ is classical, and further improvements are closely related to the distribution of zeros of the zeta-function. In particular, the best known bound for $E_k(x)$,

$$E_k(x) = O(x^{1/k} \exp(-c(k)(\log x)^{3/5}(\log \log x)^{-1/5})),$$

follows from the work of Walfisz on the error term in the Prime Number Theorem (see [32]). Still, a number of authors [1, 2, 11, 18–20, 22] have obtained sharper bounds under the assumption that the Riemann Hypothesis is true.

A related problem that has attracted considerable attention concerns the gaps between consecutive k -free integers. The first result in that direction was obtained by Fogels [10], who proved that if $\theta > 2/5$ the interval $(x, x + x^\theta]$ contains a squarefree integer for all sufficiently large x . In 1951, Roth [28] reduced the exponent $2/5$ in Fogels's result to $3/13$, while Halberstam and Roth [14] proved that the interval $(x, x + x^\theta]$ contains a k -free integer for any $\theta > 1/(2k)$ and for all sufficiently large x . Around the same time, Erdős [3] proved that there exist infinitely many intervals

$(x, x + h]$, with

$$h \gg \frac{\log x}{\log \log x},$$

which contain no squarefree integers. Together, these results inspired the conjecture that for any fixed $\varepsilon > 0$, the interval $(x, x + x^\varepsilon]$ contains a squarefree integer for sufficiently large x . This conjecture seems beyond the reach of current methods, though Granville [13] has shown that, like many other famous theorems and conjectures in number theory, it follows from the *abc*-conjecture of Masser and Oesterlé.

Initially, further improvements on Roth's result [28] on gaps between squarefree numbers were obtained through the method of exponential sums [12, 25, 26, 29], while the (mostly elementary) work of Halberstam and Roth [14] inspired research on the distribution of k -free numbers in polynomial sequences: see [15, 16, 24] for some early work and [6, §2] for a more detailed history. Starting in the late 1980s, Filaseta and Trifonov published a series of papers [4, 5, 7–9, 30, 31], where they developed an elementary proof [8] that there exists a constant $c > 0$ such that the interval $(x, x + cx^{1/5} \log x]$ contains a squarefree integer for all sufficiently large x . Later, Trifonov [31] generalized this result and proved that, for each $k \geq 3$, there exists a constant $c = c(k) > 0$ such that the interval $(x, x + cx^{1/(2k+1)} \log x]$ contains a k -free integer for all sufficiently large x . Filaseta and Trifonov [9] generalized their method to achieve progress in other problems—see the survey article [6] for the history of such developments, but sharper bounds on the gaps between k -free integers have remained elusive.

During the past couple of decades, number theorists' interest in numerically explicit results has increased significantly, and this has led to the development of numerically explicit versions of known theorems. As the Filaseta–Trifonov approach to gaps between k -free integers is both self-contained and “numerically friendly,” it therefore makes sense to investigate fully explicit versions of the results of [8] and [31]. In this note, we prove such explicit versions of the gap results for squarefree integers. Our main theorem is as follows.

Theorem 1. *For any $x \geq 2$, the interval $(x, x + 11x^{1/5} \log x]$ contains a squarefree integer.*

The reader familiar with the work of Filaseta and Trifonov may wonder whether the techniques from this work can be extended to obtain similar results on gaps between k -free integers when $k \geq 3$. This is very much possible. Indeed, we have proved the following result on gaps between cubefree integers. Its proof and the proofs of companion results on k -free integers with $k \geq 4$ will appear in a forthcoming paper.

Theorem 2. *For any $x \geq 2$, the interval $(x, x + 5x^{1/7} \log x]$ contains a cubefree integer.*

The focus of the above theorems is on providing explicit intervals that work for all x . The price we pay for this universality are the somewhat elevated values of the constants 11 and 5 in the theorems. If one is interested in reducing those constants further and willing to accept a result that holds only for sufficiently large x , then one may prefer versions like those given in the next theorem.

Theorem 3. *Every interval*

- $(x, x + 5x^{1/5} \log x]$ contains a squarefree number for $x \geq e^{400}$;
- $(x, x + 2x^{1/5} \log x]$ contains a squarefree number for $x \geq e^{1800}$;
- $(x, x + x^{1/5} \log x]$ contains a squarefree number for $x \geq e^{500000}$.

Mossinghoff, Oliveira e Silva and Trudgian [23] (see also Marmet [21]) investigated long gaps between squarefree numbers numerically. Their computational work establishes the size of the longest gaps up to 10^{18} , which are all dramatically smaller than the bounds that we get in this paper. The largest gap that they find is a string of 18 consecutive non-squarefree numbers, the first of which is 125 781 000 834 058 568. As a result of their work, we can assume $x \geq 10^{18} > e^{41}$ throughout the rest of this paper.

Theorem 3 already hints that the constants in Theorems 1 and 2 are influenced by the “small” values of x . Indeed, we establish Theorem 1 for $x \geq e^{116}$. To bridge the gap between this lower bound and e^{41} , we prove several propositions giving results with larger exponents, which are however superior to the results of the main theorem for small x . In particular, we find that the interval $(x, x + 5x^{1/4}]$ always contains a squarefree integer (Proposition 2) and the interval $(x, x + 3.8x^{1/4}]$ contains a squarefree integer for $x \geq e^{109}$ (Proposition 3).

It should be clear by now from the above discussion, that the values of the constants and the various cutoffs in the theorems (and in Propositions 2 and 3) are not exact, but rather “nice” approximations. We say more about this in Section 7.¹

Notation. Throughout the paper, for a real number θ , we use $\lfloor \theta \rfloor$ to denote the greatest integer less than or equal to θ ; also, $\{\theta\} = \theta - \lfloor \theta \rfloor$. We write $|A|$ for the size of the set A , and $\pi(x)$ for the prime counting function.

2. PRELIMINARIES

2.1. Outline of the method. Let $N(x, h)$ be the number of integers in $(x, x + h]$ that are not squarefree. Clearly, to prove any of our theorems, it suffices to show that $N(x, h) < h - 1$ for the respective choices of x and h . We first sieve this interval of the squares of very small primes, up to a parameter J to be chosen later.

¹The interested reader can explore these phenomena further using the SageMath code for the computational part of our work, which is available at <https://github.com/agreatnate/explicit-k-free-integer-bounds>

The number of integers in $(x, x + h]$ divisible by the square of a prime up to J is at most

$$h \left(1 - \prod_{p \leq J} \left(1 - \frac{1}{p^2} \right) \right) + 2^{\pi(J)} = h \left(1 - \prod_{p \leq J} \left(1 - \frac{1}{p^2} \right) + \frac{2^{\pi(J)}}{h} \right) =: h\sigma'_0(h, J).$$

We then count separately the integers divisible by p^2 for each prime $p > J$. We find that

$$N(x, h) \leq h\sigma'_0(h, J) + \sum_{p > J} \left(\left\lfloor \frac{x+h}{p^2} \right\rfloor - \left\lfloor \frac{x}{p^2} \right\rfloor \right), \quad (2.1)$$

where the sum on the right is over all primes greater than J . To bound the latter sum, we study separately the contributions of “small” and “large” primes p . We introduce a parameter H , which we will later choose as $H = mh$, with $m \geq 1$ of moderate size, and we use this parameter to split the sum in (2.1) as follows:

$$\left(\sum_{J < p \leq H} + \sum_{p > H} \right) \left(\left\lfloor \frac{x+h}{p^2} \right\rfloor - \left\lfloor \frac{x}{p^2} \right\rfloor \right) =: \Sigma_1 + \Sigma_2. \quad (2.2)$$

The contribution of the small primes can be bounded easily. We have

$$\begin{aligned} \Sigma_1 &\leq \sum_{J < p \leq H} \left(\frac{h}{p^2} + 1 \right) \leq h \sum_{p > J} \frac{1}{p^2} + \pi(H) \\ &< h \left(\sigma_1 - \sum_{p \leq J} \frac{1}{p^2} \right) + \pi(H), \end{aligned} \quad (2.3)$$

where σ_1 , the sum of the reciprocals of all squares of primes, satisfies

$$\sigma_1 < 0.4523. \quad (2.4)$$

We group the sum over primes up to J appearing in (2.3) with $\sigma'_0(h, J)$ to write

$$\sigma_0(h, J) = 1 - \prod_{p \leq J} \left(1 - \frac{1}{p^2} \right) - \sum_{p \leq J} \frac{1}{p^2} + \frac{2^{\pi(J)}}{h}, \quad (2.5)$$

so that we get

$$N(h, x) \leq h(\sigma_0(h, J) + \sigma_1) + \pi(H) + \Sigma_2. \quad (2.6)$$

The term $\pi(H)$ above can be bounded with the help of the following well-known result of Rosser and Schoenfeld [27, (3.2)].

Lemma 1. *For any $x > 1$, one has*

$$\pi(x) < \frac{x}{\log x} \left(1 + \frac{1.5}{\log x} \right). \quad (2.7)$$

Applying this lemma, we see that

$$\pi(H) < \sigma_2(h, m)h, \quad \sigma_2(h, m) := \frac{m}{\log(mh)} \left(1 + \frac{1.5}{\log(mh)} \right). \quad (2.8)$$

The estimation of the sum Σ_2 occupies the remainder of the paper. We remark that primes $p > \sqrt{2x}$ do not contribute to that sum, since for such primes we have

$$0 < \frac{x}{p^2} < \frac{x+h}{p^2} \leq \frac{2x}{p^2} < 1.$$

Moreover, if $p > h^{1/2}$, we get

$$0 \leq \left\lfloor \frac{x+h}{p^2} \right\rfloor - \left\lfloor \frac{x}{p^2} \right\rfloor \leq \frac{h}{p^2} + 1 < 2.$$

Thus, the finite sum Σ_2 counts the primes $p \in (H, \sqrt{2x}]$ for which there exists an integer m with

$$\frac{x}{p^2} < m \leq \frac{x+h}{p^2}.$$

The latter inequality can be expressed in terms of the fractional part of xp^{-2} : it says that $\{xp^{-2}\} > 1 - hp^{-2}$. Therefore,

$$\Sigma_2 \leq |S(H, \sqrt{2x})|, \quad (2.9)$$

where

$$S(M, N) := \left\{ u \in \mathbb{Z} : M < u \leq N, \gcd(u, 2) = 1, 1 - \frac{h}{u^2} \leq \left\{ \frac{x}{u^2} \right\} < 1 \right\}. \quad (2.10)$$

We remark that while we no longer require the elements of $S(M, N)$ to be prime, we do restrict them to odd values so that the differences between any two elements of the set are even, a fact which will be useful later.

Thus, in view of (2.6), (2.8) and (2.9), to prove any of our results, it will suffice to find a choice of H such that

$$|S(H, \sqrt{2x})| \leq h\sigma_3(h, m), \quad (2.11)$$

for some bounded function $\sigma_3(h, m)$ such that

$$\sigma_0(h, J) + \sigma_1 + \sigma_2(h, m) + \sigma_3(h, m) < 1 - \frac{1}{h}. \quad (2.12)$$

In Section 6, we establish inequalities of the form (2.11) and optimize the choices of several parameters to ensure that the respective versions of (2.12) hold. We conclude the present section with the statements of a couple of general-purpose lemmas, which we will use repeatedly in the remainder of the paper to obtain bounds on the spacing and cardinality of sets $S(M, N)$.

2.2. Some general lemmas. Our bounds on $|S(M, N)|$ are based on the simple idea that if the minimum distance between distinct elements of a set of integers A is at least d , then

$$|A \cap (M, N]| \leq d^{-1}(N - M) + 1. \quad (2.13)$$

In Sections 3–5, we prove several results on the spacing between elements of sets

$$S(M) := S(M, \lambda M),$$

where $\lambda > 1$ is a constant. Those spacing estimates and inequality (2.13) yield bounds on $|S(M)|$, which we leverage with the help of the next lemma.

Lemma 2. *Suppose that A_1, A_2, A_3, b_1, b_2 are positive reals and u, v, λ are real numbers with $0 < u < v < 1 < \lambda$. Assume that for all $M \in [x^u, x^v]$ the estimate*

$$|S(M)| \leq A_1 M^{b_1} + A_2 M^{-b_2} + A_3$$

holds. Then

$$|S(x^u, x^v)| \leq A'_1 x^{b_1 v} + A'_2 x^{-b_2 u} + A'_3 \log x + A_3,$$

where

$$A'_1 = \frac{A_1}{1 - \lambda^{-b_1}}, \quad A'_2 = \frac{A_2}{1 - \lambda^{-b_2}}, \quad A'_3 = A_3 \cdot \frac{v - u}{\log \lambda}.$$

Proof. This is standard: we cover the interval $(x^u, x^v]$ with intervals of the form $(M, \lambda M]$, apply the hypothesis to each of them, and sum the ensuing geometric progressions. The only (minimal) novelty in the present version is the explicit description of the coefficients A'_j in terms of the A_j 's and the various parameters. The reader will find a detailed proof of a variant for $\lambda = 2$ in [5, Lemma 1]. \square

Some of our results also rely on the properties of divided differences. For a function $f : [a, b] \rightarrow \mathbb{R}$ and $s + 1$ points $t_0, t_1, \dots, t_s \in [a, b]$, the *divided difference (of order s)*, $f[t_0, t_1, \dots, t_s]$, of f at the given points is defined recursively: we set $f[t_0] = f(t_0)$ when $s = 0$, and

$$f[t_0, t_1, \dots, t_s] = \frac{f[t_1, \dots, t_s] - f[t_0, \dots, t_{s-1}]}{t_s - t_0}$$

when $s \geq 1$. Divided differences are a tool in numerical analysis that has a long and rich history, but here we are interested only in two of their elementary properties, which we summarize in the next lemma. The reader can find proofs of these properties in many texts on numerical analysis that discuss interpolation theory: e.g., [17, Ch. 6].

Lemma 3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a function, let $t_0 < t_1 < \dots < t_s$ be distinct numbers in $[a, b]$, and let $f[t_0, t_1, \dots, t_s]$ denote the respective divided difference of f . Then*

$$f[t_0, t_1, \dots, t_s] = \sum_{j=0}^s \frac{f(t_j)}{\prod_{i \neq j} (t_j - t_i)},$$

where the product is over $i \in \{0, 1, \dots, s\} \setminus \{j\}$. Moreover, if f has s continuous derivatives on $[a, b]$, then there is a number $\xi \in (t_0, t_s)$ such that

$$f[t_0, t_1, \dots, t_s] = \frac{f^{(s)}(\xi)}{s!}.$$

3. BASIC SPACING LEMMAS

Let M be a large parameter, with $H \leq M \leq \sqrt{2x}$, and let $\lambda \in (1, 2]$ be a constant. In this section, we prove several lower bounds on the minimum distance between distinct elements of $S(M)$. As we pointed out in the introduction, the computational work in [23] allows us to assume that x is large. Also, while in our proofs we will utilize several different choices for h and H , we will always have $h \leq H$ and $h \leq 2x^{1/3}$. Thus, we assume in the remainder of the paper that

$$x \geq e^{41}, \quad 1000 \leq h \leq 2x^{1/3}. \quad (3.1)$$

3.1. Spacing for pairs. First, we show that two distinct elements of $S(M)$ cannot be “too close” to one another.

Lemma 4. *Suppose that $H \leq M$. If u and $u + a$ are distinct elements of $S(M)$, then*

$$a > 0.4995x^{-1}M^3. \quad (3.2)$$

Proof. Consider the function $f(u) = xu^{-2}$. If $u, u + a \in S(M)$, we have

$$f(u) = n_1 - \theta_1, \quad f(u + a) = n_2 - \theta_2, \quad (3.3)$$

with $n_1, n_2 \in \mathbb{Z}$, $0 < \theta_1, \theta_2 < hM^{-2}$. So,

$$f(u + a) - f(u) = n - \theta, \quad |\theta| < hM^{-2}.$$

By the mean-value theorem, there exists a number $\xi \in (u, u + a)$ such that

$$|f(u + a) - f(u)| = a|f'(\xi)| = \frac{2ax}{\xi^3} > \frac{2x}{(\lambda M)^3}.$$

If $n = 0$, we have $|f(u + a) - f(u)| = |\theta| < hM^{-2}$, and we deduce that

$$2\lambda^{-3}x < hM < 3x^{5/6},$$

which contradicts (3.1). Thus, we have $n \neq 0$, so $|n| \geq 1$. We also get that

$$|\theta| < hM^{-2} \leq hH^{-2} \leq H^{-1} \leq 0.001.$$

Hence, $|f(u + a) - f(u)| \geq 1 - |\theta| \geq 0.999$, and we obtain

$$0.999 \leq |f(u + a) - f(u)| = 2ax\xi^{-3} < 2axM^{-3},$$

from which (3.2) follows. \square

Applying (2.13) to the result of the last lemma, we obtain the following bound on the size of $S(M)$.

Corollary 1. *Under the hypotheses of Lemma 4, we have*

$$|S(M)| \leq 0.4995^{-1}(\lambda - 1)xM^{-2} + 1.$$

3.2. Spacing for triples. Next, we consider any three distinct elements $u, u + a, u + b$ of $S(M)$, with $0 < a < b$, and obtain lower bounds on b .

Lemma 5. *Let $\lambda \leq 1.2$, $m \geq 1.5$, and suppose that $mh = H \leq M$. If $0 < a < b$ and $u, u + a, u + b$ are elements of $S(M)$, then*

$$b \geq 1.3860x^{-1/3}M^{4/3}. \quad (3.4)$$

Proof. Suppose first that $b \leq 0.004M$. Write $u_1 = u$, $u_2 = u + a$, and $u_3 = u + b$, and let $n_1, n_2, n_3 \in \mathbb{Z}$ be such that

$$f(u_i) = n_i - \theta_i, \quad 0 < \theta_i < hM^{-2} \quad (i = 1, 2, 3).$$

We consider the second divided difference $f[u_1, u_2, u_3]$. By Lemma 3,

$$\begin{aligned} f[u_1, u_2, u_3] &= \frac{f(u_1)(u_3 - u_2) + f(u_2)(u_1 - u_3) + f(u_3)(u_2 - u_1)}{(u_2 - u_1)(u_3 - u_1)(u_3 - u_2)} \\ &= \frac{(n_1 - \theta_1)(b - a) - (n_2 - \theta_2)b + (n_3 - \theta_3)a}{ab(b - a)} =: \frac{n - \theta}{V}, \end{aligned}$$

where

$$n = (b - a)n_1 - bn_2 + an_3 \quad \text{and} \quad \theta = (b - a)\theta_1 - b\theta_2 + a\theta_3.$$

In particular, since $\theta_i > 0$, we have

$$-bhM^{-2} < -b\theta_2 < \theta < (b - a)\theta_1 + a\theta_3 < bhM^{-2}.$$

Moreover, since $u, u + a$ and $u + b$ are all odd (see (2.10)) we know that a and b are both even, so n must be as well.

We will show that $n \neq 0$. Suppose that $n = 0$. Then

$$|f[u_1, u_2, u_3]| = \frac{|\theta|}{V} < \frac{bhM^{-2}}{ab(b - a)} = \frac{h}{a(b - a)M^2} < \frac{hx}{0.999M^5},$$

after an appeal to (3.2) and the bound $b - a \geq 2$. However, using Lemma 3, we also get that

$$|f[u_1, u_2, u_3]| = \frac{|f''(\xi)|}{2!} = \frac{3x}{\xi^4} \geq \frac{3x}{(\lambda M)^4}.$$

Thus,

$$\frac{3x}{(\lambda M)^4} < \frac{hx}{0.999M^5} < \frac{1.002hx}{HM^4},$$

which contradicts the hypotheses of the lemma.

Having proved that $n \neq 0$ and using that it is even, we find that $|n| \geq 2$. Hence,

$$|f[u_1, u_2, u_3]| = \frac{|n - \theta|}{V} \geq \frac{2 - |\theta|}{ab(b - a)} > \frac{1.997}{ab(b - a)}, \quad (3.5)$$

since

$$|\theta| < bhM^{-2} < 0.004hH^{-1} \leq \frac{1}{250m} \leq \frac{1}{375}.$$

On the other hand, by Lemma 3,

$$|f[u_1, u_2, u_3]| = \frac{3x}{\xi^4} \leq \frac{3x}{M^4}. \quad (3.6)$$

From (3.5), (3.6), and the elementary inequality $a(b-a) \leq \frac{1}{4}b^2$, we deduce that

$$\frac{3b^3x}{4} \geq 3ab(b-a)x > 1.997M^4, \quad (3.7)$$

and the conclusion of the lemma follows in the case $b \leq 0.004M$.

Finally, when $b > 0.004M$, we have

$$b^3 > (0.004M)^3 > \frac{8}{3}x^{-1}M^4,$$

by the assumptions that $M \leq \sqrt{2x}$ and $x \geq e^{41}$. \square

Note that the expression on the right side of (3.4) is a lower bound for the minimum distances between successive elements of the set $S_1(M)$ containing every other element of $S(M)$. Since $|S(M)| \leq 2|S_1(M)|$, this observation and (2.13) yields the following corollary.

Corollary 2. *Under the hypotheses of Lemma 5, we have*

$$|S(M)| \leq 1.4430(\lambda - 1)x^{1/3}M^{-1/3} + 2.$$

4. SPACING FOR PAIRS OF PAIRS

In this section, we study a special family of quadruples $u, u+a, u+b, u+a+b$ of elements of $S(M)$. The special form of the spacing between the four numbers allows us to obtain bounds on b that are stronger than those for general quadruples in $S(M)$; in the next section, we will average these bounds over b . In the next lemma, we use the third-order divided difference of $f(u) = xu^{-2}$ for the points $u, u+a, u+b$, and $u+a+b$ to bound b from below.

Lemma 6. *Let $\lambda \leq 1.05$, $m \geq 5$, and suppose that $mh \leq H \leq M$. If $0 < a < 2a \leq b$ and $u, u+a, u+b, u+a+b$ are elements of $S(M)$, then*

$$ab^3 \geq 0.6600x^{-1}M^5. \quad (4.1)$$

Proof. Consider points $u_1 = u, u_2 = u+a, u_3 = u+b$, and $u_4 = u+a+b$ in $S(M)$. Recall that by the definition of the set $S(M)$, there exist integers n_1, \dots, n_4 and reals $\theta_1, \dots, \theta_4$ such that

$$f(u_i) = n_i - \theta_i, \quad 0 < \theta_i < hM^{-2} \quad (1 \leq i \leq 4). \quad (4.2)$$

We consider the divided difference $f[u_1, \dots, u_4]$.

Due to the special configuration of the distances between the four points, the formula in Lemma 3 simplifies to

$$f[u_1, u_2, u_3, u_4] = \frac{f(u_4) - f(u_1)}{ab(a+b)} - \frac{f(u_3) - f(u_2)}{ab(b-a)} =: \frac{n - \theta}{V},$$

where $V = ab(a+b)(b-a)$ and

$$\begin{aligned} n &= (b-a)(n_4 - n_1) - (a+b)(n_3 - n_2), \\ \theta &= (b-a)(\theta_4 - \theta_1) - (a+b)(\theta_3 - \theta_2). \end{aligned}$$

We remark that n is an even integer and $|\theta| < 2bhM^{-2}$.

We will show that $n \neq 0$. Suppose that $n = 0$. Then

$$|f[u_1, \dots, u_4]| = \frac{|-\theta|}{V} \leq \frac{2bhM^{-2}}{ab(a+b)(b-a)}.$$

Recalling (3.7), we deduce that

$$|f[u_1, \dots, u_4]| < \frac{2hM^{-2}}{ab(b-a)} \leq \frac{6hx}{1.997M^6}.$$

However, Lemma 3 gives

$$|f[u_1, \dots, u_4]| = \frac{|f^{(3)}(\xi)|}{3!} = \frac{4x}{\xi^5} \geq \frac{4x}{(\lambda M)^5},$$

for some $\xi \in (M, \lambda M]$. We combine these upper and lower bounds to get

$$\frac{4x}{(\lambda M)^5} < \frac{6hx}{1.997M^6} < \frac{3.005hx}{HM^5},$$

which contradicts the assumptions of the lemma.

Since n is even and nonzero, we can now use that $|n| \geq 2$ combined with the observation $b^2 - a^2 \geq 0.75b^2$ to obtain

$$|f[u_1, \dots, u_4]| = \frac{|n - \theta|}{V} \geq \frac{2 - |\theta|}{ab(b^2 - a^2)} \geq \frac{1.98}{0.75ab^3}, \quad (4.3)$$

since

$$|\theta| < 2bhM^{-2} < 2(\lambda - 1)hH^{-1} \leq \frac{1}{10m} \leq \frac{1}{50}.$$

On the other hand, by Lemma 3,

$$|f[u_1, \dots, u_4]| = \frac{4x}{\xi^5} \leq \frac{4x}{M^5}. \quad (4.4)$$

The lemma follows from (4.3) and (4.4). \square

Our next result is of a somewhat different nature from the spacing lemmas established hitherto. In this lemma, instead of proving that the distance b between the two pairs exceeds some lower bound in terms of x , M , and possibly, a , we establish

a kind of a dichotomy for b : either $b \geq B_1$ for some lower bound B_1 , or $b \leq B_2$, with B_2 significantly smaller than B_1 .

Lemma 7. *Let $\lambda \leq 1.05$, $m \geq 5$, and suppose that $mh \leq H \leq M$. If $0 < a < 2a \leq b$ and $u, u+a, u+b, u+a+b$ are elements of $S(M)$, then exactly one of the conditions*

$$a^3b < \lambda^6hx^{-1}M^4, \quad (4.5)$$

or

$$a^3b > (0.5 - \lambda m^{-1})x^{-1}M^5, \quad (4.6)$$

must hold.

Proof. We start from the algebraic identity

$$\frac{2u+3a}{(u+a)^2} - \frac{2u-a}{u^2} = \frac{a^3}{u^2(u+a)^2}.$$

Since $u, u+a \in S(M)$, we can use this identity and (4.2) to get that

$$\frac{a^3x}{u^2(u+a)^2} = \frac{(2u+3a)x}{(u+a)^2} - \frac{(2u-a)x}{u^2} = n' + \theta' \quad (4.7)$$

where $n' = (2u+3a)n_2 - (2u-a)n_1$ is an even integer and

$$|\theta'| = |\theta_1(2u-a) - \theta_2(2u+3a)| \leq 2u|\theta_1 - \theta_2| + a(\theta_1 + 3\theta_2) < (2u+4a)hM^{-2}.$$

Combining (4.7) with the analogous identity for the pair $u+b, u+a+b$, we find that

$$\frac{a^3x}{u^2(u+a)^2} - \frac{a^3x}{(u+b)^2(u+a+b)^2} = n + \theta, \quad (4.8)$$

where $n \in \mathbb{Z}$ is even and

$$|\theta| < (4u+2b+8a)hM^{-2} \leq 4(u+a+b)hM^{-2} \leq 4\lambda hM^{-1} \leq 4\lambda m^{-1}.$$

Next we observe, by the mean-value theorem, there is a $\xi \in (u, u+b)$ such that

$$\frac{a^3x}{u^2(u+a)^2} - \frac{a^3x}{(u+b)^2(u+a+b)^2} = \frac{2a^3bx(2\xi+a)}{\xi^3(\xi+a)^3}.$$

This expression is bounded above by

$$\frac{2a^3bx(2\xi+a)}{\xi^3(\xi+a)^3} < \frac{4a^3bx}{\xi^3(\xi+a)^2} < 4a^3bxM^{-5}, \quad (4.9)$$

and bounded below by

$$\frac{2a^3bx(2\xi+a)}{\xi^3(\xi+a)^3} > \frac{4a^3bx}{\xi^2(\xi+a)^3} > 4a^3bx(\lambda M)^{-5}. \quad (4.10)$$

When $a^3b \leq (0.5 - \lambda m^{-1})x^{-1}M^5$, (4.8), (4.9), and the bound on $|\theta|$ yield

$$n - 4\lambda m^{-1} < n + \theta < 4a^3bxM^{-5} \leq 2 - 4\lambda m^{-1},$$

and hence, $n < 2$. On the other hand, if $a^3b \geq \lambda^6hx^{-1}M^4$, we deduce from (4.8) and (4.10) that

$$4\lambda hM^{-1} < 4a^3bx(\lambda M)^{-5} < n + \theta < n + 4\lambda hM^{-1},$$

so in this case $n > 0$. Since n is an even integer, it can satisfy only one of the conditions $n > 0$ and $n < 2$; therefore, at least one of (4.5) or (4.6) must hold. This completes the proof, since under the hypotheses of the lemma, the lower bound in (4.6) exceeds the upper bound in (4.5) at least by a constant factor. \square

5. THE MAIN BOUNDS ON $|S(M)|$

Let

$$A = 1.3860x^{-1/3}M^{4/3}. \quad (5.1)$$

In Section 3, we proved that $b \geq A$ whenever $u, u+a, u+b$ are distinct elements of $S(M)$. Therefore, if u_0, u_1, \dots, u_s are the elements of $S(M)$, listed in increasing order, the set $S_1(M) = \{u_0, u_2, u_4, \dots\}$ has no gaps $< A$ and satisfies

$$|S(M)| \leq 2|S_1(M)|. \quad (5.2)$$

In this section, we use (5.2) and the lemmas in the last section to prove the following result.

Proposition 1. *Suppose $h = 11x^{1/5} \log x$, let $\lambda = 1.045$ and $x \geq e^{116}$, and suppose that $5.5h \leq M \leq x^{2/5}$. Then*

$$|S(M)| \leq h(\sigma_3(M) + \sigma_4(M)), \quad (5.3)$$

where

$$\sigma_3(M) = (0.5298x^{1/5} + 0.3400x^{-1/5}M)h^{-1} + 0.0308x^{1/15}M^{-1/3}, \quad (5.4)$$

and

$$\sigma_4(M) = \begin{cases} 1.2105x^{-2/3}M^{7/3} & \text{if } M \leq 5x^{1/4}, \\ 1.4182x^{-1/9}M^{1/9} & \text{if } M > 5x^{1/4}. \end{cases} \quad (5.5)$$

Remark 1. Notice that when x is relatively small, the condition $M \leq 5x^{1/4}$ in Proposition 1 is impossible, and so only the second condition will be used in the range of “small” values of x .

The proof of this proposition uses the set

$$T(M; a) = \{u : u, u+a \text{ are consecutive elements of } S_1(M)\}$$

to bound $|S_1(M)|$. The starting point is the elementary identity

$$|S_1(M)| = 1 + \sum_{a=1}^{\infty} |T(M; a)| = 1 + \sum_{a \geq A} |T(M; a)|, \quad (5.6)$$

which is a direct consequence of the definition of $T(M; a)$. Further, for any $B \geq A$, we have

$$\sum_{a \geq B} a|T(M; a)| \leq \sum_{a \geq A} a|T(M; a)| \leq (\lambda - 1)M + 1,$$

so

$$\sum_{a \geq B} |T(M; a)| \leq (\lambda - 1)MB^{-1} + B^{-1}.$$

Applying this inequality to the right side of (5.6), we find, for any parameter $B \geq 2$, that

$$|S_1(M)| \leq 1.5 + (\lambda - 1)MB^{-1} + \sum_{A \leq a < B} |T(M; a)|. \quad (5.7)$$

5.1. Proof of Proposition 1. We recall the quantity A defined in (5.1), and we select

$$B = \delta x^{-1/5}M, \quad \delta = 0.17, \quad (5.8)$$

in the imminent application of (5.7). We fix an integer a , with $A \leq a \leq B$. If u_0, u_1, \dots, u_t are the elements of $T(M; a)$, listed in increasing order, the set $T_1(M) = \{u_0, u_2, u_4, \dots\}$ contains only elements of $T(M; a)$ such that if $u, u + b \in T_1(M)$, then $b \geq 2a$. Clearly, $|T(M; a)| \leq 2|T_1(M)|$.

Let I be a subinterval of $(M, \lambda M]$ of length

$$|I| = (0.5 - \lambda m^{-1})a^{-3}x^{-1}M^5,$$

and let $u, u + b$ be two elements of $T_1(M) \cap I$. Since $b \geq 2a$, we can apply Lemma 7 to show that b must satisfy (4.5). Taking u and $u + b$ to be the smallest and largest elements of $T_1(M) \cap I$ respectively, we can use this bound on b to deduce that the set $T_1(M) \cap I$ is contained in an interval of length $\leq \lambda^6 a^{-3} h x^{-1} M^4$. Furthermore, by (4.1), we have that

$$b \geq 0.8706a^{-1/3}x^{-1/3}M^{5/3}.$$

Combining these two observations we find that

$$|T_1(M) \cap I| \leq \frac{\lambda^6 a^{-3} h x^{-1} M^4}{0.8706a^{-1/3}x^{-1/3}M^{5/3}} + 1 < 1.4959a^{-8/3}h x^{-2/3}M^{7/3} + 1. \quad (5.9)$$

Since we need at most

$$\frac{(\lambda - 1)M}{(0.5 - \lambda m^{-1})a^{-3}x^{-1}M^5} + 1 < 0.1452a^3 x M^{-4} + 1 \quad (5.10)$$

intervals of length $|I|$ to cover $(M, \lambda M]$, we conclude that

$$\begin{aligned} |T_1(M)| &\leq (0.1452a^3 x M^{-4} + 1) (1.4959a^{-8/3}h x^{-2/3}M^{7/3} + 1) \\ &< 0.2173a^{1/3}h x^{1/3}M^{-5/3} + 0.1452a^3 x M^{-4} + 1.4959a^{-8/3}h x^{-2/3}M^{7/3} + 1. \end{aligned}$$

Thus,

$$|T(M; a)| < 0.4346a^{1/3}hx^{1/3}M^{-5/3} + 0.2904a^3xM^{-4} \\ + 2.9918a^{-8/3}hx^{-2/3}M^{7/3} + 2. \quad (5.11)$$

Next, we use (5.11) to bound the right side of (5.7). With our choice of parameters, (5.7) gives

$$|S_1(M)| \leq 1.5 + 0.045\delta^{-1}x^{1/5} + \sum_{A \leq a < B} |T(M; a)|. \quad (5.12)$$

Thus, we need to sum each of the four terms on the right side of (5.11) over $a \in [A, B)$. Recalling the inequality

$$\sum_{k \leq K} k^s < \frac{(K+1)^{s+1}}{s+1} \quad (s > 0),$$

and noting that $B = \delta Mx^{-1/5} \geq 5.5h\delta x^{-1/5} > 10.285 \log x > 1193$, we find that

$$\sum_{\substack{2 \leq a \leq B \\ a \text{ even}}} a^s < \frac{(B+2)^{s+1}}{2(s+1)} < \frac{(1.002B)^{s+1}}{2(s+1)}. \quad (5.13)$$

Hence,

$$0.4346hx^{1/3}M^{-5/3} \sum_{\substack{A \leq a < B \\ a \text{ even}}} a^{1/3} < \frac{0.4346 \cdot (1.002B)^{4/3}}{8/3} hx^{1/3}M^{-5/3} \\ < 0.0154hx^{1/15}M^{-1/3}, \quad (5.14)$$

and

$$0.2904xM^{-4} \sum_{\substack{A \leq a < B \\ a \text{ even}}} a^3 < \frac{0.2904 \cdot (1.002B)^4}{8} xM^{-4} < 0.00004x^{1/5}. \quad (5.15)$$

Combining (5.8), (5.11), (5.12), (5.14), and (5.15), we conclude that

$$|S_1(M)| \leq h\sigma'_3(M) + 2.9918hx^{-2/3}M^{7/3} \sum_{\substack{A \leq a < B \\ a \text{ even}}} a^{-8/3}, \quad (5.16)$$

where

$$\sigma'_3(M) = (0.2649x^{1/5} + 0.17x^{-1/5}M)h^{-1} + 0.0154x^{1/15}M^{-1/3}. \quad (5.17)$$

We estimate the sum on the right side of (5.16) in different ways, depending on the size of M . When $M \leq 5x^{1/4}$, we use that

$$\sum_{\substack{A \leq a < B \\ a \text{ even}}} a^{-8/3} < \frac{\zeta(8/3)}{2^{8/3}} < 0.2023. \quad (5.18)$$

On the other hand, when $M > 5x^{1/4}$, we have $A > 1.386 \cdot 5^{4/3} > 11.8501$, so

$$\sum_{\substack{A \leq a < B \\ a \text{ even}}} a^{-8/3} < \frac{3}{5 \cdot 2^{8/3}} \left(\frac{A}{2} - 1 \right)^{-5/3} < 0.4083A^{-5/3} < 0.237x^{5/9}M^{-20/9}. \quad (5.19)$$

The proposition follows from (5.2) and (5.16)–(5.19). \square

6. PROOF OF THEOREM 1

The proof of the theorem uses different approaches for different values of x . As we stated in the introduction, the work in [23] establishes our result (and much more) for $x \leq e^{41}$. In Section 6.1, we focus on large x and show that for $x \geq e^{116}$, Theorem 1 follows from Proposition 1. To complete the proof, in Section 6.2, we prove two asymptotically weaker variants, which are, however, stronger than the theorem for small x . Those alternative results establish Theorem 1 in the intermediate range $e^{41} \leq x \leq e^{116}$.

6.1. Large x . Let $x \geq e^{116}$ and set $H = 5.5h$ in (2.2) and (2.11). First, we use Proposition 1 and Lemma 2 to bound $|S(H, x^{2/5})|$.

Suppose first that $H \leq 5x^{1/4}$, (in this case we can assume $x \geq e^{150}$) we split $S(H, x^{2/5})$ in two pieces to account for the different cases in (5.5). When we apply Lemma 2 to the bound (5.3) for $M \in [H, 5x^{1/4}]$, we find that

$$\begin{aligned} |S(H, 5x^{1/4})| &< h \left(\frac{1.2105 \cdot 5^{7/3} x^{-1/12}}{1 - 1.045^{-7/3}} + \frac{0.0308 x^{1/15} H^{-1/3}}{1 - 1.045^{-1/3}} \right) + \frac{1.7x^{1/20}}{1 - 1.045^{-1}} \\ &\quad + 0.5298x^{1/5} \left(\frac{\log(5x^{1/4}/H)}{\log(1.045)} + 1 \right) \\ &< 0.1034h + 0.0001x^{1/5} + 0.5298x^{1/5} \left(\frac{\log x}{20 \log(1.045)} - \frac{\log(12.1 \log x)}{\log(1.045)} + 1 \right) \\ &< 0.1034h + 0.6019x^{1/5} \log x - 89.7886x^{1/5} < 0.1582h. \end{aligned}$$

Similarly, when we apply Lemma 2 to (5.3) for $M \in [5x^{1/4}, x^{2/5}]$, we get

$$\begin{aligned} |S(5x^{1/4}, x^{2/5})| &< h \left(\frac{1.4182x^{-1/15}}{1 - 1.045^{-1/9}} + \frac{0.0308 \cdot 5^{-1/3} x^{-1/60}}{1 - 1.045^{-1/3}} \right) + \frac{0.34x^{1/5}}{1 - 1.045^{-1}} \\ &\quad + 0.5298x^{1/5} \left(\frac{3 \log x}{20 \log(1.045)} - \frac{\log 5}{\log(1.045)} + 1 \right) \\ &< 0.1148h + 1.8055x^{1/5} \log x - 10.9463x^{1/5} < 0.2790h. \end{aligned}$$

Hence,

$$|S(H, x^{2/5})| < 0.1582h + 0.2790h = 0.4372h. \quad (6.1)$$

Next, we consider the case $H > 5x^{1/4}$ (which implies that $x \leq e^{151}$). In this case we need only consider the latter case of Proposition 1 for M in the full range $(H, x^{2/5}]$. Applying Lemma 2 in this situation gives

$$\begin{aligned} |S(H, x^{2/5})| &< h \left(\frac{1.4182x^{-1/15}}{1 - 1.045^{-1/9}} + \frac{0.0308x^{1/15}H^{-1/3}}{1 - 1.045^{-1/3}} \right) + \frac{0.34x^{1/5}}{1 - 1.045^{-1}} \\ &\quad + 0.5298x^{1/5} \left(\frac{\log x}{5 \log(1.045)} - \frac{\log(60.5 \log x)}{\log(1.045)} + 1 \right) \\ &< 0.2378h + 2.4073x^{1/5} \log x - 98.1700x^{1/5} < 0.3976h, \end{aligned} \quad (6.2)$$

on noting that $98.170x^{1/5} > 0.059h$ when $x \leq e^{151}$.

To complete the estimation of $|S(H, \sqrt{2x})|$, we apply Lemma 2 to the bound in Corollary 1 for $M \in [x^{2/5}, \sqrt{2x}]$. This yields

$$|S(x^{2/5}, \sqrt{2x})| < \frac{0.0901x^{1/5}}{1 - 1.045^{-2}} + \frac{\log x}{10 \log(1.045)} + \frac{0.5 \log 2}{\log(1.045)} + 1 < 0.0009h. \quad (6.3)$$

Together, (6.1)–(6.3) establish (2.11) with

$$\sigma_3 = \begin{cases} 0.4381 & \text{if } H \leq 5x^{1/4}, \\ 0.3985 & \text{if } H > 5x^{1/4}, \end{cases}$$

for all $x \geq e^{116}$. Taking $J = 120$ in (2.5), we have $\sigma_0(h, 120) \leq -0.0595$ in the same range. Furthermore, for all $x \geq e^{116}$, we have $\sigma_2(h, 5.5) < 0.1797$, and for $x \geq e^{150}$, we have $\sigma_2(h, 5.5) < 0.1461$. Thus,

$$\sigma_0(h, 120) + \sigma_1 + \sigma_2(h, 5.5) + \sigma_3(h, 5.5) < \begin{cases} 0.9770 & \text{if } H \leq 5x^{1/4}, \\ 0.9710 & \text{if } H > 5x^{1/4}, \end{cases}$$

which establishes (2.12), and therefore the theorem, for $x \geq e^{116}$.

6.2. Intermediate x . Suppose that $x \geq e^{41}$. We consider $h = 5x^{1/4}$ and choose $\lambda = 1.025$, $J = 19$ and $H = 1.75h$. With these choices, we apply Lemma 2 to the result of Corollary 2 to obtain

$$\begin{aligned} |S(H, \sqrt{2x})| &< \frac{1.4430 \cdot (0.025)x^{1/3}H^{-1/3}}{1 - 1.025^{-1/3}} + 2 \left(\frac{\log(\sqrt{2x}) - \log H}{\log(1.025)} + 1 \right) \\ &< 0.4272h + \frac{\log x}{2 \log(1.025)} + \frac{\log 2 - 2 \log(8.75/1.025)}{\log(1.025)} < 0.4331h. \end{aligned}$$

That is, (2.12) holds with $\sigma_3(h, 1.75) = 0.4331$. Moreover, when $h = 5x^{1/4}$ and $x \geq e^{41}$, we have

$$\sigma_0(h, 19) \leq -0.0543, \quad \sigma_2(h, 1.75) \leq 0.158.$$

Thus, when $h = 5x^{1/4}$ and $x \geq e^{41}$, we have

$$\sigma_0(h, 19) + \sigma_1 + \sigma_2(h, 1.75) + \sigma_3(h, 1.75) < 0.9891.$$

Together with the computations of [23], this proves the following result.

Proposition 2. *For any $x \geq 2$, the interval $(x, x + 5x^{1/4}]$ contains a squarefree integer.*

Moreover, an identical calculation for $x \geq e^{109}$ with $h = 3.8x^{1/4}$, $H = 4.5h$, $\lambda = 1.0001$, and $J = 100$ yields

$$\begin{aligned} \sigma_0(h, 100) + \sigma_1 + \sigma_2(h, 4.5) + \sigma_3(h, 4.5) \\ < -0.0594 + 0.4523 + 0.1571 + 0.4423 = 0.9924, \end{aligned}$$

which yields the following alternative.

Proposition 3. *For any $x \geq e^{109}$, the interval $(x, x + 3.8x^{1/4}]$ contains a squarefree integer.*

Since $5x^{1/4} \leq 11x^{1/5} \log x$ for $x \leq e^{109.7}$, Proposition 2 implies Theorem 1 for $x \leq e^{109}$. Finally, since $3.8x^{1/4} \leq 11x^{1/5} \log x$ for $x \leq e^{116.3}$, Proposition 3 establishes Theorem 1 when $e^{109} \leq x \leq e^{116}$. This completes the proof of the theorem.

7. ASYMPTOTIC RESULTS AND FINAL COMMENTS

We conclude by noting a few of the explicit bounds that can be obtained by these methods if one no longer requires the bounds to be admissible for all values of $x \geq 2$, allowing instead results valid for sufficiently large values of x .

Some of the possible results that can be obtained by tweaking the parameters used in the proof of Theorem 1 are given in the statement of Theorem 3. To prove any of those results, we reset the parameters m, J, λ, δ that appear in the proofs of Proposition 1 and Theorem 1 and then update the various constants. (When x is as large as in Theorem 3, the inequality $H \leq 5x^{1/4}$ always holds, so only the first case in the proof of Theorem 1 can occur.) To establish the claims of Theorem 3, we always select $J = 100$, $\lambda = 1.02$, and $m = \sqrt{\log x_0}$, where x_0 is the lower bound on x in each result; we only vary the choice of δ . For example, when $h = 5x^{1/5} \log x$, $x \geq e^{400}$ (hence, $m = 20$), and $\delta = 0.3$, we have

$$\sigma_0(h, 100) + \sigma_1 + \sigma_2(h, m) + \sigma_3(h, m) < 0.9811.$$

For $h = 2x^{1/5} \log x$ and $x \geq e^{1800}$, the choice $\delta = 0.6$ yields an upper bound of 0.9857; and for $h = x^{1/5} \log x$ and $x \geq e^{500000}$, $\delta = 0.87$ gives a bound of 0.9981.

Remark 2. Looking back at the proofs of our theorems, one can see that the value of h in our theorems is of the form $h(x) = cx^{1/5} \log x$, with c an upper bound for a rather complicated bounded function $C(x; m, J, \lambda, \delta)$, which is decreasing in the variable x . Once x is sufficiently large, the decay in x appears to overwhelm the effect of the other parameters. On the other hand, to claim a specific value

of c for all $x \geq x_0$, one generally needs to find acceptable choice of the other parameters to ensure that (2.12) holds. It seems that if one were to make the function $C(x; m, J, \lambda, \delta)$ fully explicit, one may even be able to identify a four-dimensional neighborhood of the chosen values of m, J, λ, δ such that all the choices of the parameters in that neighborhood are acceptable.

Acknowledgments. This work is the result of an REU project that took place on the campus of Towson University during the summer of 2022, with the financial support of the National Science Foundation under grants DMS-2136890 and DMS-2149865. The authors also acknowledge financial support from the Fisher College of Science and Mathematics and TU's Mathematics Department. We also thank M. Filaseta, O. Trifonov, and T. Trudgian for commenting on early drafts of this work and to the anonymous referee for their careful reading of the manuscript and for making an important suggestion regarding the overall organization of the paper.

REFERENCES

- [1] R. C. Baker and J. Pintz, *The distribution of squarefree numbers*, Acta Arith. **46** (1985), no. 1, 73–79. MR831264
- [2] R. C. Baker and K. Powell, *The distribution of k -free numbers*, Acta Math. Hungar. **126** (2010), no. 1-2, 181–197. MR2593323
- [3] P. Erdős, *Some problems and results in elementary number theory*, Publ. Math. Debrecen **2** (1951), 103–109. MR45759
- [4] M. Filaseta, *An elementary approach to short interval results for k -free numbers*, J. Number Theory **30** (1988), no. 2, 208–225. MR961917
- [5] ———, *Short interval results for squarefree numbers*, J. Number Theory **35** (1990), no. 2, 128–149. MR1057318
- [6] M. Filaseta, S. W. Graham, and O. Trifonov, *Starting with gaps between k -free numbers*, Int. J. Number Theory **11** (2015), no. 5, 1411–1435. MR3376218
- [7] M. Filaseta and O. Trifonov, *On gaps between squarefree numbers*, Analytic Number Theory (Allerton Park, IL, 1989), Progr. Math., vol. 85, Birkhäuser Boston, Boston, MA, 1990, pp. 235–253. MR1084183
- [8] ———, *On gaps between squarefree numbers. II*, J. London Math. Soc. (2) **45** (1992), no. 2, 215–221. MR1171549
- [9] ———, *The distribution of fractional parts with applications to gap results in number theory*, Proc. London Math. Soc. (3) **73** (1996), no. 2, 241–278. MR1397690
- [10] E. Fogels, *On average values of arithmetic functions*, Proc. Cambridge Philos. Soc. **37** (1941), 358–372. MR4843
- [11] S. W. Graham, *The distribution of squarefree numbers*, J. London Math. Soc. (2) **24** (1981), no. 1, 54–64. MR623670
- [12] S. W. Graham and G. Kolesnik, *On the difference between consecutive squarefree integers*, Acta Arith. **49** (1988), no. 5, 435–447. MR967330
- [13] A. Granville, *ABC allows us to count squarefrees*, Internat. Math. Res. Notices (1998), no. 19, 991–1009. MR1654759
- [14] H. Halberstam and K. F. Roth, *On the gaps between consecutive k -free integers*, J. London Math. Soc. **26** (1951), 268–273. MR43120
- [15] C. Hooley, *On the power free values of polynomials*, Mathematika **14** (1967), 21–26. MR214556

- [16] M. N. Huxley and M. Nair, *Power free values of polynomials. III*, Proc. London Math. Soc. (3) **41** (1980), no. 1, 66–82. MR579717
- [17] E. Isaacson and H. B. Keller, *Analysis of Numerical Methods*, Dover Publications, Inc., New York, 1994, Corrected reprint of the 1966 original. MR1280462
- [18] C.-H. Jia, *The distribution of square-free numbers*, Sci. China Ser. A **36** (1993), no. 2, 154–169. MR1223084
- [19] H.-Q. Liu, *On the distribution of k -free integers*, Acta Math. Hungar. **144** (2014), no. 2, 269–284. MR3274401
- [20] ———, *On the distribution of squarefree numbers*, J. Number Theory **159** (2016), 202–222. MR3412720
- [21] L. Marmet, *First occurrences of square-free gaps and an algorithm for their computation*, preprint arXiv:1210.3829, 2012.
- [22] H. L. Montgomery and R. C. Vaughan, *The distribution of squarefree numbers*, Recent Progress in Analytic Number Theory, Vol. 1 (Durham, 1979), Academic Press, London-New York, 1981, pp. 247–256. MR637350
- [23] Michael J. Mossinghoff, Tomás Oliveira e Silva, and Timothy S. Trudgian, *The distribution of k -free numbers*, Math. Comp. **90** (2021), no. 328, 907–929. MR4194167
- [24] M. Nair, *Power free values of polynomials. II*, Proc. London Math. Soc. (3) **38** (1979), no. 2, 353–368. MR531167
- [25] R. A. Rankin, *Van der Corput's method and the theory of exponent pairs*, Quart. J. Math. Oxford Ser. (2) **6** (1955), 147–153. MR72170
- [26] H.-E. Richert, *On the difference between consecutive squarefree numbers*, J. London Math. Soc. **29** (1954), 16–20. MR57898
- [27] J. B. Rosser and L. Schoenfeld, *Approximate formulas for some functions of prime numbers*, Illinois J. Math. **6** (1962), 64–94. MR137689
- [28] K. F. Roth, *On the gaps between squarefree numbers*, J. London Math. Soc. **26** (1951), 263–268. MR43119
- [29] P. G. Schmidt, *Abschätzungen bei unsymmetrischen Gitterpunktproblemen*, Dissertation, Göttingen, 1964, Dissertation zur Erlangung des Doktorgrades der Mathematisch-Naturwissenschaftlichen Fakultät der Georg-August-Universität zu Göttingen. MR0181609
- [30] O. Trifonov, *On the squarefree problem. II*, Math. Balkanica (N.S.) **3** (1989), no. 3-4, 284–295. MR1048051
- [31] ———, *On gaps between k -free numbers*, J. Number Theory **55** (1995), no. 1, 46–59. MR1361558
- [32] A. Walfisz, *Weylsche Exponentialsummen in der neueren Zahlentheorie*, Mathematische Forschungsberichte, XV, VEB Deutscher Verlag der Wissenschaften, Berlin, 1963. MR0220685

DEPARTMENT OF MATHEMATICS, TOWSON UNIVERSITY, TOWSON, MD 21252, U.S.A.
Email address: `akumchev@towson.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA BERKELEY, BERKELEY, CA 94720, U.S.A.
Email address: `wademc@berkeley.edu`

DEPARTMENT OF MATHEMATICS, TOWSON UNIVERSITY, TOWSON, MD 21252, U.S.A.
Email address: `nmcnew@towson.edu`

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, U.S.A.
Email address: `arianap@mit.edu`

DEPARTMENT OF MATHEMATICS, TOWSON UNIVERSITY, TOWSON, MD 21252, U.S.A.
Email address: `rscher4@students.towson.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA BERKELEY, BERKELEY, CA 94720, U.S.A.
Email address: `simon_ziehr@berkeley.edu`